

EO 4 Ecosystem Accounting 2022



Customized Ecosystem Service Models for Crop and Carbon Stock by ARIES Platform

K.HAYASHI Nagoya University
H.Okazawa Tokyo University of Agriculture
F. Villa BC3
S. Balbi BC3
K. Zhang Tokyo University of Agriculture

Date EO4EA 30 NOV, 2022



→ THE EUROPEAN SPACE AGENCY



ARtificial Intelligence for Ecosystem Services

ARIES is a networked software technology that redefines ecosystem service assessment and valuation for decision-making. The ARIES approach to mapping natural capital, natural processes, human beneficiaries, and service flows to society is a powerful new way to visualize, value, and manage the ecosystems on which the human economy and well-being depend.

[Learn more](#)

k.LAB Japan

Nagoya University Prof. Hayashi
Osaka University Dr. machimura
Chubu University Dr. Sugita
Tokyo University of Agriculture Prof. Okazawa
NIES
etc.




ARIES(Artificial Intelligence for Environment and Sustainability)

#ARIES15years:
Celebrating 15 years of AI for environmental sustainability

Since 2007, ARIES has been building the first "Wikipedia-like" open-source platform for interoperable data and models, through AI. Check out this infographic providing an overview of our vision to take open science to the next level and watch for a series of new ARIES online infographics to be released throughout the following months!


[Learn more](#)




Three+ one ARIES/k.LAB interface




k.Explorer




ARIES-for-SEEA explorer



k.Modeler



k.LAB Japan model



→ THE EUROPEAN SPACE AGENCY

ARIES Global ES model and k.LAB Japan ES model



Carbon stock: G-carbon, J-Carbon stock

Climate regulation: J-Forest volume, J-LST

Air regulation: J-NO2, J-SO2, etc.

Flood regulation: G-flood regulation, *J-Flood Luration area*

Pollination: G-pollination

Landslide prevention: G-landside, G-sediment, *J-Sediment Disaster hazard area*

Biomass: G-Biomass(Maize crop, wood), J-crop model, J-tree biomass

Water: G-water(under development)

Culture: G-Recreation, *National park, Nature conservation area, Wildlife area, Local tourism resources, Urban park*

MAIN EO data used:

- ALOS by JAXA: images
- HRLULC by JAXA: LULC map
- AW3D by JAXA, NTT data, RESTEC: Height data

Simplified Crop model based on AquaCrop by FAO



Landcover type: Agriculture land

ESA. Land Cover CCI Product User Guide Version 2. Tech. Rep. (2017). Available at: maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf

HRLULC(Japan)

HRLULC: High-Resolution Land Use and Land Cover Map by JAXA, https://www.eorc.jaxa.jp/ALOS/en/dataset/lulc_e.htm

Relation between canopy cover and NDVI

ZHANG et.al.(2021) IJERD,12-2

Evapotranspiration

Antonio Trabucco, Robert J. Zomer, Global Aridity Index and Potential Evapo-Transpiration (ET0) Climate Database v2

This study is supported by Kakenhi 20K06351, JSPS, Japan.


Parameters:

Several AquaCrop parameters

Harvest rate, Leftover ratio

Output


- Biomass
- Yield
- Leftover



Relation between HRLULC(Japan) and forest canopy(LiDAR and AW3D)volume and carbon stock(Field survey)

LiDAR data(air plane), AW3D

- DSM(digital surface model):
 - LiDAR 1-2m-grid, AW3D 2.5-5m, 30m
- HRLULC: 10m-grid



EO challenges, opportunities and recommendations



k.LAB Japan model

EO is useful for ES assessment.

Processed data: Small grid scale:
10m, 2.5m~

- Height data
- Land use and land cover data
- Image, NDVI, Evapotranspiration

Combination with field measurement
results